

Morphometric Evaluation of T₁₃-L₁ and L₁-L₂ Intervertebral Space for Subarachnoid Injection in Simmental and Brown Swiss Cattle: A Cadaver Study

Yalçın AKBULUT¹ , Özgür AKSOY² , Sadık YAYLA² , Savaş ÖZTÜRK² , Uğur AYDIN² , İsa ÖZAYDIN² 

¹Department of Anatomy, Kafkas University, Faculty of Medicine, Kars, Turkey

²Department of Surgery, Kafkas University, Faculty of Veterinary Medicine, Kars, Turkey

Cite this article as: Akbulut, Y., Aksoy, Ö., Yayla, S., Öztürk, S., Aydın, U., & Özyayın, İ. (2021). Morphometric Evaluation of T₁₃-L₁ and L₁-L₂ Intervertebral Space for Subarachnoid Injection in Simmental and Brown Swiss Cattle: A Cadaver Study. *Acta Veterinaria Eurasia*, 47(2), 93-97.

ORCID IDs of the authors: Y.A. 0000-0003-4661-2224; Ö.A. 0000-0002-4800-6079; S.Y. 0000-0001-6734-421X; S.Ö. 0000-0001-6934-4176; U.A. 0000-0001-5756-4841; İ.Ö. 0000-0003-4652-6377.

Abstract

This study aimed to reveal the probable anatomic differences of the T₁₃-L₁ or L₁-L₂ vertebrae in Simmental and Brown Swiss cattle during surgeries, such as subarachnoid anesthesia and cerebrospinal fluid aspiration, and thus determine a reference value. This study was conducted in 30 cows, of which 15 were Brown Swiss cattle and 15 were Simmental, aged 5–6 years, and having a live weight of 400–450 kg. The region from the T₁₃-L₁ vertebrae to the sacrum was taken from each animal and divided into 2 equal parts throughout the long axis. The distance between the spinous processes of the T₁₃-L₁ and L₁-L₂ vertebrae, spinous processes length of the T₁₃ and L₁ vertebrae,

sagittal length of interarcuate spaces between the T₁₃-L₁ and L₁-L₂ vertebrae, and vertical length of vertebral canals of the T₁₃ and L₁ vertebrae were measured using an electronic digital caliper. Statistical results showed that although there was no significant difference within the groups ($p > .05$), there was a significant difference between the groups in terms of the sagittal length of interarcuate space between the T₁₃-L₁ vertebra ($p < .05$). We believe that this result is because of the anatomical and genetic differences between the 2 cattle breeds.

Keywords: Cattle, interarcuate space, morphometry

Introduction

Intrathecal anesthesia is preferred nowadays for paralumbar, perineal, posterior extremities, and other surgeries in cattle owing to the adverse effects of general anesthesia. Examination of the cerebrospinal fluid (CSF) can be done via CSF aspiration through the subarachnoid entrance using radio-diagnostic procedures (Kılıç et al., 2015; Yayla et al., 2012; Yayla & Kılıç, 2010). Therefore, acknowledging the anatomic structure of the region according to the breed will contribute to more effective anesthesia and other procedures.

All vertebrae, except the atlas, involve the vertebral body and vertebral arch in mammals. The 2 neighboring vertebrae are connected to each other from their vertebral body through

the intervertebral disc and ventral and dorsal longitudinal ligaments. In addition, neighboring vertebrae are connected to each other from their vertebral arch by means of the interarcuate ligament, intertransverse ligament, interspinous ligament, and supraspinous ligament. The vertebral arch and vertebral body are merged through the pedicle of the vertebral arch. The lamina of the vertebral arch, an enlarged part, is located on the pedicle of the vertebral arch. The 2 laminae of the vertebral arch of a vertebra unite in a median line to form the spinous process (Dursun, 2000; König & Liebich, 2007; Sharshar et al., 2015).

Although the atlanto-occipital, lumbosacral, or thoracolumbar space is preferred for CSF aspiration in cattle, only the lumbosacral space is used for spinal anesthesia (Aksoy et al., 2012;

The pre-findings of the study were presented orally in Congress of national veterinary surgery organized in Antalya, Turkey, between 23 and 26 October 2014.

Address for Correspondence: Yalçın AKBULUT • **E-mail:** yalcinakbulut@kafkas.edu.tr

Received Date: January 4, 2021 • **Accepted Date:** March 5, 2021 • DOI: 10.5152/actavet.2021.21001

Available online at actavet.org



This work is licensed under a Creative Commons Attribution-NonCommercial 4.0 International License.

De Lahunta & Glass, 2009; Hiraoka et al., 2007; Lee et al., 2004; Skarda et al., 1989; Vandeveldel et al., 2001). However, thoracolumbar injection is not recommended in the surgery of the paralumbar fossa or flank, and Cesarean, rumenotomy, and abomasopexy are preferably conducted in the standing position; therefore, subarachnoid injection should be done through the T₁₃-L₁ vertebrae or alternatively through the L₁₋₂ vertebrae (Caulkett et al., 1993; De Rossi et al., 2003; Lee & Yamada, 2005; Seyrek-İntaş et al., 2001; Yayla et al., 2013).

In a previous study (Aksoy et al., 2012), subarachnoid entries were performed through the T₁₃-L₁ or L₁₋₂ vertebrae in different animal breeds by the same doctor, and it was detected that this procedure was more difficult in the Simmental breed than in other breeds. In addition, the fact that the clinicians at our university had difficulty entering the subarachnoid space of Simmental breed cattle brought to our hospital, but could do so easily in Brown Swiss breeds encouraged us to perform this study. This problem also bring about the question of whether the spaces T₁₃-L₁ or L₁₋₂ will show morphometric changes according to different breed or not. This study aimed to find answers to these questions also measure the intervertebral space distances between T₁₃-L₁ and L₁₋₂ vertebrae in Simmental and Brown Swiss cattle, to determine whether there is any difference between the two cattle breeds, and thus to establish a reference for surgical operations such as subarachnoid anesthesia or CSF aspiration.

Method

The use of the animals in this study was allowed by the local ethics committee of animal experiments of Kafkas University (KAU-HADYEK/2015-025/08.01.2015). This investigation was performed in 30 cows, of which 15 were Brown Swiss cattle (group I) and 15 were Simmental (group II). They were aged 5–6 years, with 400–450 kg live weight, did not have any disorder or trauma complaint, and were ethically slaughtered in the Kars slaughter house. The vertebral column, with its soft tissues, till the tail vertebra and T₁₃ of each cow was excised (Figure 1, 2) and divided into 2 equal parts and moved to the anatomy laboratory of the Kafkas University. Following this, the distance between the spinous processes of the T₁₃ and L₁ vertebrae (Figure 1 “a”), distance between the spinous processes of the L₁-L₂ vertebrae (Figure 1 “b”), lengths of spinous processes of the T₁₃

(Figure 1 “c”) and L₁ vertebrae (Figure 1 “d”), sagittal length of the intercaruate spaces between the T₁₃-L₁ (Figure 1 “e”) and L₁-L₂ (Figure 1 “f”), and vertical lengths of the vertebral canals of the T₁₃ (Figure 1 “g”) and L₁ vertebrae (Figure 1 “h”) were measured using an electronic digital caliper (.01 mm, BTS, England). The obtained data were analyzed statistically using Minitab packaged software (Trialware, Pennsylvania, USA) and then evaluated within groups and between groups comparatively using one-way analysis of variance (Tukey’s test). To compare variables between the groups, t test was used.

Results

According to the results of the study, there was no statistically significant difference within the groups in terms of the distance between the spinous processes of the T₁₃ and L₁ vertebrae, distance between the spinous processes of the L₁-L₂ vertebrae, lengths of spinous processes of the T₁₃ and L₁ vertebrae, sagittal length of the intercaruate spaces between the T₁₃-L₁ and L₁-L₂ vertebrae, and vertical lengths of the vertebral canals of the T₁₃ and L₁ vertebrae (p > .05) (Figure 1). However, a significant difference was found between the groups in terms of the sagittal length of intercaruate space between the T₁₃-L₁ (p < .05). There was no statistically significant difference within the groups (Table 1).



Figure 1

Sagittal Section of the Thoracolumbar and Os Sacrum in Brown Swiss Cattle. (a) Distance between the Spinous Processes of T₁₃-L₁ Vertebrae, (b) Distance between the Spinous Processes of L₁₋₂ Vertebrae, (c) The Spinous Process Length of T₁₃ Vertebra, (d) Length of Spinous Processes of L₁ Vertebra, (e) Sagittal Length of Intercaruate space between T₁₃-L₁ Vertebrae, (f) Sagittal Length of Intercaruate Space Between L₁₋₂ Vertebrae, (g) Vertical Length of Vertebral Canal T₁₃ Vertebra, (h) Vertical Length of Vertebral Canal L₁ Vertebra.

Table 1

Comparison between the Groups of Morphometric Measurement Results Obtained from the Study

Groups	Distance between the spinous processes (mm)		Spinous processes length (mm)		Sagittal length of intercaruate space (mm)		Vertical length of vertebral canal (mm)	
	T ₁₃ -L ₁	L ₁ -L ₂	T ₁₃	L ₁	T ₁₃ -L ₁	L ₁ -L ₂	T ₁₃	L ₁
(n = 15)	19.17 ± 5.72	18.59 ± 5.14	59.88 ± 5.07	54.63 ± 5.48	5.12 ± 1.20	5.47 ± 1.35	18.78 ± 2.28	19.82 ± 2.49
(n = 15)	18.86 ± 3.08	19.13 ± 4.22	60.15 ± 7.49	55.59 ± 6.12	4.15 ± 1.0*	4.79 ± 1.34	19.07 ± 1.26	19.82 ± 1.91

Note. *: Statistical significance between groups in each column (p < .05). Values are presented as mean ± standard deviation




Figure 2

Sagittal Section of the Thoracolumbar and Os Sacrum in Simmental Cattle. (a) Distance between the Spinous Processes of T_{13} - L_1 Vertebrae, (b) Distance between the Spinous Processes of L_{1-2} Vertebrae, (c) The Spinous Process Length of T_{13} Vertebra, (d) Length of Spinous Processes of L_1 Vertebra, (e) Sagittal Length of Intercaruate Space between T_{13} - L_1 Vertebrae, (f) Sagittal Length of Intercaruate Space between L_{1-2} Vertebrae, (g) Vertical Length of Vertebral Canal T_{13} Vertebra, (h) Vertical Length of Vertebral Canal L_1 Vertebra.

Discussion, and Conclusion and Recommendations

Subarachnoid or intrathecal administration is necessary primarily for intrathecal anesthesia, radio-diagnostic procedures, and CSF aspiration. Although lumbosacral space is favored for intrathecal anesthesia in many animal breeds, cisternal or lumbosacral injections are used for myelography (Aksoy et al., 2012; Yayla et al., 2012). Many surgeries need to be performed in the standing position because of the known complications of surgeries, especially in adult cattle (Aksoy et al., 2012; Caulkett et al., 1993; De Lahunta & Glass, 2009). Therefore, there are many studies on intrathecal or subarachnoid anesthesia and analgesia, which do not let the animals lie down (Aksoy et al., 2012; Yayla & Kiliç, 2010); some researchers have recommended thoracolumbar administration along with lumbosacral (Caulkett et al., 1993; Ferheller et al., 2004; Lee et al., 2006; Levis et al., 1999). Moreover, there are studies on entering via the lumbosacral line, then advancing the spinal catheter cranially or caudally after having entered through the thoracolumbar space for the blockage of nerves in the L_{2-5} level, and administering the anesthetic drug to this part (Aksoy et al., 2012; Özaydın & Kiliç, 2003; Skarda et al., 1989). These studies have also reported that T_{13} - L_1 intrathecal entrance was difficult; thus, L_{1-2} may be an alternative for T_{13} - L_1 injection. However, studies on morphometric measurements of the mentioned regions for subarachnoid introduction in cattle are limited. In a previous study (Aksoy et al., 2012), subarachnoid analgesia was performed in different cattle breeds, but the subarachnoid injection in Simmental cows was determined to be problematic. This prompted us to conduct this study, and the intervertebral distance was measured between the T_{13} - L_1 and L_{1-2} vertebrae of Simmental and Brown Swiss cattle (the 2 common breeds in cultural breeding). We aimed to determine a reference value for surgical operations, such as subarachnoid anesthesia or CSF aspiration, from this region. We believe that the results of this study will contribute to the literature by providing further information on subarachnoid surgeries.

Although there are no studies conducted on morphometric measurements in cattle, several studies reported the distance till the subarachnoid gap on the skin during intrathecal attempts. Lee et al. (2004) found the distance between the skin and subarachnoid space to be 84–93 mm in 8 cows and 85–95 mm in 10 cows; they did not find any statistically significant difference in the effect on analgesia in these 2 groups. Another study (Lee et al., 2006) found this distance to be 82–91 mm in adult animals and 57–65 mm in young animals, whereas another study (Lee et al., 2004) found this to be 89 mm. Hiraoka et al. (2007) reported this distance to be 81–90 mm. Our study dealt with the length of spinous processes, except for the skin. This length was determined as 59.88 ± 5.07 mm for T_{13} vertebra in Brown Swiss cattle and 60.15 ± 7.49 mm in Simmentals, whereas it was found to be 54.63 ± 5.48 mm for L_1 vertebra in Brown Swiss cattle and 55.59 ± 6.12 mm in Simmentals. In addition, there was no statistically significant difference both within the group and between group measurements.

Sinding and Berg (2010) carried out a study dealing with similar data in warmblood foals, and no interspinous space width less than 4 mm was found. The width between T_{16-17} was determined to be 5.9 ± 1.2 mm, and the widest interspinous space was reported as 8.9 ± 2.6 mm between T_{10-12} . However, the researchers did not conduct any study in cattle to investigate the spinous processes of T_{13} - L_1 and L_{1-2} vertebrae in terms of morphometric values. In the present study, the distance between the spinous processes of T_{13} - L_1 vertebrae was determined as 19.17 ± 5.72 mm in Brown Swiss cattle and 18.86 ± 3.08 mm in Simmentals; the distance between the spinous processes of L_{1-2} vertebrae was 18.59 ± 5.14 mm in Brown Swiss cattle and 19.13 ± 4.22 mm in Simmentals. Furthermore, there was no statistically significant difference in measurements both within the group and between the groups. Therefore, it can be assumed that these 2 breeds have substantial similarities.

In a study carried out in water buffalo by using computed tomography (Sharshar et al., 2015), the intercaruate space be-

tween the last thoracic spine and initial lumbar vertebra and within all the lumbar vertebrae were determined, although not between the last 2 thoracic spines. However, there was no study regarding the sagittal length of intercaruate space in cattle. Therefore, to the best of our knowledge, this is the first study investigating the sagittal length of intercaruate space in Brown Swiss and Simmental cattle. This length was found to be 5.12 ± 1.20 mm in Brown Swiss cattle and 4.15 ± 1.0 mm in Simmental in $T_{13}-L_1$, whereas it was 5.47 ± 1.35 mm in Brown Swiss cattle and 4.79 ± 1.34 mm in Simmental in L_{1-2} . There was no statistically significant difference in these data within the groups. Although there was no statistically significant difference in the sagittal length of L_{1-2} intercaruate space, a statistically significant difference was determined for $T_{13}-L_1$ ($p < .05$). Therefore, we assume that $T_{13}-L_1$ intrathecal injection would be easier for Brown Swiss cattle than Simmental ones.

The diameter of spinal canal of L_1 vertebra was specified as 16.89, 8.83, and 11.81 mm for human, sheep, and deer, respectively (Bai et al., 2012). The diameter of the spinal canal of T_{13} vertebra was found to be 9.4 and 9.8 mm for L_1 vertebra (Mageed et al., 2013). This study revealed that the vertical length of the vertebral canal of T_{13} vertebra was 18.78 ± 2.28 mm in Brown Swiss cattle and 19.07 ± 1.26 mm in Simmentals, whereas the vertical length of the vertebral canal of L_1 vertebra was 19.82 ± 2.49 mm in Brown Swiss cattle and 19.82 ± 1.91 mm in Simmentals. Furthermore, they were statistically similar both within the group and between groups.

In conclusion, this study was designed to compare the intervertebral space distances between the $T_{13}-L_1$ and L_{1-2} vertebrae in Brown Swiss cattle and Simmentals. There may be difficulties in subarachnoid entry in Simmentals as the sagittal length of $T_{13}-L_1$ intercaruate space and sagittal length of intercaruate space between L_{1-2} vertebrae is narrower than Brown Swiss cattle. We think that this difference is the result of the Simmental cattle breed genetically being more muscular and having higher milk production.

Ethics Committee Approval: The use of the animals in this study was allowed by the local ethics committee of animal experiments of Kafkas University (KAU-HADYEK/2015-025/08.01.2015).

Peer-review: Externally peer-reviewed.

Author Contributions: Concept - İ.Ö., Ö.A.; Design - S.Y., Y.A.; Supervision - Ö.A., İ.Ö.; Resources - U.A., S.Ö.; Materials - Y.A., U.A.; Data Collection and/or Processing - Y.A., U.A.; Analysis and/or Interpretation - S.Y., Y.A., U.A.; Literature Search - Ö.A., S.Y., Y.A.; Writing Manuscript - Y.A., S.Y.; Critical Review - İ.Ö., S.Y., S.Ö.

Conflict of Interest: The authors have no conflicts of interest to declare.

Financial Disclosure: This study was financially supported by the scientific research project committee of the Kafkas University (KAU-BAP, 2014-TS-32).

References

- Aksoy, Ö., Özaydin, İ., Kırmızıgül, A. H., Kılıç, E., Öztürk, S., Kurt, B., Yayla, S., Sözmen, M., & Atakışi, E. (2012). Evaluation of experimental subarachnoid analgesia with a combination of detomidine and ropivacaine for flank analgesia in cows. *Veterinarski Arhiv*, 82(5), 463-472.
- Bai, X., Liu, G., Xu, C., Zhuang, Y., & Zhang, J. (2012). Morphometry research of deer, sheep, and human lumbar spine: Feasibility of using deer and sheep in spinal animal models. *International Journal of Morphology*, 30(2), 510-520. [Crossref]
- Caulkett, N., Cribb, P. H., & Duke, T. (1993). Xylazine epidural analgesia for caesarean section in cattle. *Canadian Veterinary Journal*, 34(11), 674-676.
- De Lahunta, A., & Glass, E. (2009). Cerebrospinal fluid and hydrocephalus. In *Veterinary Neuroanatomy and Clinical Neurology* (pp: 54-76). Saunders Elsevier, St Louis. [Crossref]
- De Rossi, R., Gaspar, E. B., Junqueira, A. L., & Beretta, M. P. (2003). A comparison of two subarachnoid α_2 -agonists, xylazine and clonidine, with respect to duration of antinociception, and hemodynamic effects in goats. *Small Ruminant Research*, 47(2), 103-111. [Crossref]
- Dursun, N. (2000). *Veterinary Anatomy II* (1st ed.) Medisan Press, Ankara, Turkey.
- Ferheller, E. E., Caulkett, N. G., & Balley, J. V. (2004). A romifidine and morphine combination for epidural analgesia of the flank in cattle. *Canadian Veterinary Journal*, 45(11), 917-923.
- Hiraoka, M., Miyagawa, T., Kobayashi, H., Takahashi, T., & Kishi, H. (2007). Successful introduction of modified dorsolumbar epidural anaesthesia in a bovine referral center. *Journal of Veterinary Science*, 8(2), 181-184. [Crossref]
- Kılıç, E., Yayla, S., Kamiloğlu, A., Baran, V., & Öğün, M. (2015). Effects of intrathecal administration of ketamine HCl in young calves: a clinical trial. *Bull Vet Inst Pulawy*, 59(1), 155-159. [Crossref]
- Konig, H. E., & Liebich, H. G. (2007). *Veterinary Anatomy of Domestic Mammals, Textbook and Colour Atlas* (3rd ed.) Schattauer GmbH, Stuttgart, Germany.
- Lee, I., & Yamada, H. (2005). Epidural administration of fixed volumes of xylazine and lidocaine for anaesthesia of dairy cattle undergoing flank surgery. *Journal of the American Veterinary Medical Association*, 227(5), 781-784. [Crossref]
- Lee, I., Yamagishi, N., Oboshi, K., Sasaki, N., & Yamada, H. (2004). Comparison of xylazine, lidocaine and the two drugs combined for modified dorsolumbar epidural anaesthesia in cattle. *Veterinary Record*, 155(25), 797-799.
- Lee, I., Yamagishi, N., Oboshi, K., Sasaki, N., & Yamada, H. (2006). Practical tips for modified dorsolumbar epidural anaesthesia in cattle. *Journal of Veterinary Science*, 7(1), 69-72. [Crossref]
- Levis, C. A., Constable, P. D., Huhn, J. C., & Morin, D. E. (1999). Sedation with xylazine and lumbosacral epidural administration of lidocaine and xylazine for umbilical surgery in calves. *Journal of the American Veterinary Medical Association*, 214(1), 89-95.
- Mageed, M., Berner, D., Julke, H., Hohaus, C., & Brehm, W. (2013). Morphometrical dimensions of the sheep thoracolumbar vertebrae as seen on digitised CT images. *Laboratory Animal Research*, 29(3), 138-147. [Crossref]
- Özaydin, İ., & Kılıç, E. (2003). Lumbosacral intrathecal anaesthesia with isobaric bupivacaine in cattle. *Indian Veterinary Journal*, 80(3), 540-542.
- Seyrek-İntaş, D., Topal, A., Seyrek-İntaş, K., Rocken, Kırmızıgül, A. H., & Cihan, M. (2001). Untersuchungen zum subarachnoidalen thor-

- akolumbalen Anästhesie mit Detomidin beim Pferd. *Pferdeheilkunde*, 17(1), 220-224. [\[Crossref\]](#)
- Sharshar, A., Abedellaah, B., Shoghy, K., & Rashed, R. (2015). Dorsolumbar epidural analgesia in Water Buffalo: anaesthetic assessment and anatomical studies. *Alexandria Journal of Veterinary Sciences*, 45(1), 63-70. [\[Crossref\]](#)
- Sinding, M. F., & Berg, L. C. (2010). Distances between thoracic spinous processes in Warmblood foals: a radiographic study. *Equine Veterinary Journal*, 42(6), 500-503. [\[Crossref\]](#)
- Skarda, R. T., Muir, W. W., & Hubbel, J. A. E. (1989). Comparative study of continuous lumbar segmental epidural and subarachnoid analgesia in Holstein cows. *American Journal of Veterinary Research*, 50(1), 39-44.
- Vandeveldel, M., Jaggy, A., & Lang J. (2001). Untersuchung des Liquor cerebrospinalis. In M. Vandeveldel, A. Jaggy, J. Lang (Eds.), *Veterinärmedizinische Neurologie* (pp:63-69). Berlin: Parey Verlag.
- Yayla, S., & Kılıç, E. (2010). The comparison of clinical, histopathological and some hemodynamic effects of spinal anesthesia applied in dogs through bupivacaine HCl and ropivacaine HCl in two different concentrations. *Journal of the Faculty of Veterinary Medicine Kafkas University*, 16(5), 835-840.
- Yayla, S., Kaçar, C., Kaya, D., Merhan, O., Aksoy, Ö., Kılıç, E., & Kaya, S. (2012). Clinical, biochemical and haemodynamic effects of the intrathecal ketamine for ovariohysterectomy in bitches. *Bulletin of the Veterinary Institute in Pulawy*, 56(3), 299-303. [\[Crossref\]](#)
- Yayla, S., Kılıç, E., Aksoy, Ö., Özyayın, İ., Öğün, M., & Steagall, P. V. M. (2013). The effects of subarachnoid administration of hyperbaric solutions of bupivacaine or in xylazine-sedated calves undergoing surgery. *Veterinary Record*, 173(23), 580. [\[Crossref\]](#)